

Bay-o-Pet®  
**DRONTAL®**

Allwormer chewable

for Dogs

**DISPOSAL**

Dispose of empty packaging by wrapping with paper and placing in garbage.

**STORAGE**

Store below 30°C (Room Temperature).

**APVMA Approval No 59292/52960**

**Limitation of Liability:** Failure to use the product strictly as directed may be illegal, prove dangerous or render the product ineffective. Notwithstanding anything else contained on this label, and subject to any rights you have under the Australian Consumer Law and any other applicable law, Bayer Australia Limited's liability is limited to replacing this product or supplying equivalent product or paying the cost of replacing this product, at its discretion.

BAY-O-PET®, DRONTAL® and DRONCIT® are registered trademarks of Bayer AG, Leverkusen, Germany.  
Manufactured in Australia.

For a free treatment reminder visit  
[www.friendsforever.com.au](http://www.friendsforever.com.au)

**Bayer**

BAYER AUSTRALIA LIMITED  
875 Pacific Highway  
PYMBLE N.S.W. 2073  
ACN 000 138 714



3865731

FOR ANIMAL  
TREATMENT ONLY



Bay-o-Pet®

**DRONTAL®**

for Dogs

Allwormer chewable

EACH CHEW CONTAINS: 50mg PRAZIQUANTEL, 49.8mg PYRANTEL as PYRANTEL EMBONATE, 250mg FEBANTEL

- Controls all 12 Gastrointestinal Worms & the shedding of Giardia oocysts

DOES NOT CONTROL  
HEARTWORM IN DOGS

ONE  
CHEW  
TREATS  
10KG

2 TASTY CHEWS

DRONTAL Chewable is recommended for the treatment and control of all gastrointestinal worms in dogs including:

- Roundworm *Toxocara canis*, *Toxascaris leonina*.
- Hookworm *Ancylostoma caninum*, *Ancylostoma braziliense* (this hookworm is usually restricted to tropical areas in Australia), *Ancylostoma ceylanicum*, *Uncinaria stenocephala*.
- Whipworm *Trichuris vulpis*.
- Tapeworm *Echinococcus granulosus* (hydatid tapeworm), *Dipylidium caninum* (common flea tapeworm), *Taenia* spp., *Taenia ovis*, *T. pisiformis*, *T. hydatigena*.
- and to control the shedding of oocysts of the protozoan *Giardia* sp. in dogs.

Does not control heartworm in dogs.

**DIRECTIONS FOR USE**

Administer the dog's correct dose from the hand as a treat. DRONTAL Chewable is highly palatable and 98% of dogs will consume it this way.

Where necessary for dosage the product may be cut with a sharp knife to produce half chews. If necessary, DRONTAL Chewable may be crushed and mixed in canned dog food.

It is essential to weigh dogs before treatment.

**DOSAGE**

For intestinal worms - 1 chew per 10kg bodyweight administered once.


Bodyweight	Chews
3.1-5kg	½
5.1-10kg	1
10.1-15kg	1½
15.1-20kg	2
Over 20kg	1 chew for every 10kg bodyweight.*

\*Round off doses to the nearest ½ chew above.

If worm problems persist, consult a veterinarian.

For *Giardia* infection: Administer the above dosage for three consecutive days. DRONTAL should **only** be used when a *Giardia* sp. infection has been diagnosed by a veterinarian.

To remove chew,  
punch through from front.  
Tamper evident pack.



#### HYDATID TAPEWORM

A single dose of DRONTAL Chewable is all that is required. However, this dose should be repeated every 6 weeks to ensure that newly acquired hydatid tapeworms are expelled before reaching maturity.

**Dogs should not be fed, or allowed to feed on, offal from any species. Dogs in hydatid areas should be treated every 6 weeks.**

It is important to ensure that the dog does not acquire offal in order to break the life cycle of the hydatid tapeworm.

It should be noted that hydatid tapeworm poses a severe risk to human health by transmission to humans from infected dogs.

#### HOOKWORMS & ROUNDWORMS

■ Dogs & Puppies: **Treat at 2, 4, 8 and 12 weeks** and then every month until six months of age; **thereafter every 3 months** or on veterinary advice.

■ Pregnant Bitches: **Treat at mating, then 10 days before whelping, at 2 and 4 weeks after whelping, then every 3 months.**

#### WHIPWORMS

**Treat every 6 to 8 weeks after 3 months of age.**

#### TAPEWORMS

**Treat every 3 months** except in hydatid tapeworm areas where every 6 weeks is recommended, or on veterinary advice.

*Dipylidium caninum* tapeworms are transmitted by fleas. Control of this tapeworm will require effective control of the local flea population. Worms, if present, may not appear in the droppings at all, or may not appear for up to 48hrs after treatment.

#### GIARDIA

This protozoan parasite is found commonly in dogs and may be the cause of diarrhoea, particularly in pups. It is more common amongst dogs living in multi-dog households and kennels. *Giardia* also occurs in humans and has historically been considered a potential zoonosis, however recent studies suggest that host-specific species may exist. Control of *Giardia* is difficult because the oocysts are resistant and long-lived in the environment. Diagnosis of suspected *Giardia* infection should be confirmed by a veterinary surgeon. Some evidence indicates that

bathing and a move to a non-contaminated environment in conjunction with treatment will improve overall control.

#### COMPATIBILITY

DRONTAL Chewable may be combined with other treatments e.g. heartworm prevention, flea or tick treatments.

#### SAFETY

DRONTAL Chewable may be given safely to all dogs including old animals, young pups, working dogs and greyhounds.

It may be given to pregnant bitches but is only clinically indicated in the last 10 days prior to whelping.

#### SIDE EFFECTS

Vomiting may be observed in a small number of dogs after administration.

#### RETREATMENT RECOMMENDATIONS

Worms often reinfest dogs so it is necessary to:

- treat the dog itself,
- reduce environmental contamination by daily disposal of droppings,
- remove tapeworm intermediate hosts e.g. fleas.

#### FOLLOW THE RETREATMENT RECOMMENDATIONS IN THE TABLE BELOW:

**PUPPIES** treat at 2, 4, 8, 12 weeks; then 4, 5, 6 months of age.

**ADULTS (over 6 months)** every 3 months. Every 6 weeks in Hydatid infested locations. If Whipworm is a problem retreat every 6-8 weeks.

**PREGNANT/LACTATING BITCHES** treat at mating, then 10 days before whelping and at 2 and 4 weeks after whelping, then every 3 months.

These recommendations are of a general nature.

In specific instances where infestation is particularly heavy or many dogs are housed together, a veterinary surgeon should be consulted.

Flea tapeworms can reinfest dogs and begin producing segments 2-3 weeks after a treatment.

If segments reappear before the next DRONTAL Chewable treatment is due, either retreat with DRONTAL Chewable or use DRONCIT which is a specific treatment for tapeworm.



CUSTOMER INFORMATION LINE

**1800 678 368 TOLL FREE**

From anywhere in Australia 9.00am to 4.00pm (Eastern Time) Mon-Fri

[www.friendsforever.com.au](http://www.friendsforever.com.au)